

Survey

Functional activities of the human T-cell leukemia virus type I Tax oncoprotein: cellular signaling through NF- κ B

Kuan-Teh Jeang *

Laboratory of Molecular Microbiology, National Institute of Allergy and Infectious Diseases, Room 306, Building 4, 4 Center Drive, MSC 0460, Bethesda, MD 20892-0460, USA

Abstract

Human T-cell leukemia virus type I (HTLV-I) is the etiological agent for adult T-cell leukemia (ATL), as well as for tropical spastic paraparesis (TSP) and HTLV-I associate myelopathy (HAM). A biological understanding of the involvement of HTLV-I and in ATL has focused significantly on the workings of the virally-encoded 40 kDa phospho-oncoprotein, Tax. Tax is a transcriptional activator. Its ability to modulate the expression and function of many cellular genes has been reasoned to be a major contributory mechanism explaining HTLV-I-mediated transformation of cells. In activating cellular gene expression, Tax impinges upon several cellular signal-transduction pathways, including those for CREB/ATF and NF- κ B. In this paper, we review aspects of Tax's transcriptional potential with particular focus on recent evidence linking Tax to IKK (I κ B-kinase)-complex and MAP3Ks (mitogen-activated protein kinase kinase kinases). Published by Elsevier Science Ltd.

Keywords: HTLV; Adult T-cell leukemia; Cellular transformation; NF- κ B; Tax; CREB; AP-1; HAM/TSP

Contents

| | |
|---|-----|
| 1. HTLV-I and adult T-cell leukemia (ATL) | 207 |
| 2. Nuclear transcriptional activities of Tax | 208 |
| 2.1. SRF and AP-1 | 209 |
| 2.2. CREB, CBP, P/CAF. | 209 |
| 3. Signaling by Tax through NF- κ B | 209 |
| 3.1. Interaction of Tax and NF- κ B in the nucleus | 210 |
| 3.2. Tax and cytoplasmic signaling of NF- κ B. | 210 |
| 4. Implications for cellular transformation | 213 |
| Acknowledgements | 213 |
| References | 213 |

1. HTLV-I and adult T-cell leukemia (ATL)

Human T cell leukemia virus type 1 (HTLV-I) is the etiological agent for adult T cell leukemia (ATL) [1],

* Tel.: +1-301-4966680; fax: +1-301-4803686.
E-mail address: kj7e@nih.gov (K.-T. Jeang).

tropical spastic paraparesis (TSP) and various neurological disorders termed HTLV-I-associated myelopathy (HAM) [2]. While the linkage between TSP/HAM and HTLV-I is less understood, the role of HTLV-I in ATL has been well-investigated. In this regard, *in vivo*, ATL cells show nuclei with morphological aberrations (indented or lobulated; [1]) and abnormal karyotypes [3,4]; in particular, trisomy of chromosomes 3 and 7, as well as deletions and breaks in chromosome 6 have been observed [5–8]. ATL cells frequently present as pleiomorphic multinucleated giant cells [9,10]. Consistent with *in vivo* findings, lymphocytes infected *in vitro* with HTLV-I show profound chromosomal changes [11,12] verifying a link between viral infection and morphological changes in infected cells.

While it is not yet fully understood how HTLV-I engenders ATL, several lines of evidence link the virally encoded 40 kDa nuclear phosphoprotein [13], Tax, to cellular transformation [14–19]. Thus, Tax has been shown to immortalize T lymphocytes [20] and transform rat fibroblasts [14]. Tax not only sufficiently initiates the immortalization and transformation of human thymocytes, cord blood lymphocytes and murine fibroblast cells *in vitro* [14,16,20], but in some settings has been shown to function in maintaining the transformed phenotype [21]. Tax-transformed fibroblast and lymphoid cells induced tumors *in vivo* when injected into nude mice [14,21,22]. Interestingly, overexpression of HTLV-1 Tax in transgenic mice resulted in the formation of mesenchymal tumors [23], salivary and lacrimal gland exocrinopathy [24], lymphadenopathy or splenomegaly [25] and lymphoma or leukemia [26].

Finally, when Tax is deleted from the HTLV-I genome, this altered proviral molecule is lost for its original transforming potential [27].

Mechanistically, the role of Tax in cellular transformation likely relates to its activity as a transcriptional activator. Tax potently stimulates the expression of its cognate viral LTR [28,29], as well as the promoters of several cellular genes. Thus, cellular genes such as IL-2, IL-2R, *c-fos*, GM-CSF [19,30–38], as well as β -polymerase, *c-myc*, *Lck* and p53 [39–43] have all been shown to be influenced by expression of Tax in cells.

In understanding Tax's pleiotropic transcriptional activities, work from several laboratories have collectively defined its ability to act through four discrete cellular signaling pathways: CREB/ATF [44]; NF- κ B [45]; AP-1 [46]; and SRF [47] (Fig. 1). Amongst these pathways, Tax's signaling through CREB/ATF and NF- κ B has been most extensively investigated. Here, we review in brief our current understanding of Tax's transcriptional activity, with particular emphasis on recent findings pertaining to its signaling through NF- κ B. For further discussions of these topics, readers are encouraged to consult several excellent recent reviews [45,48,49].

2. Nuclear transcriptional activities of Tax

The HTLV-I Tax protein is predominantly a viral nuclear antigen [50] with a well-defined nuclear localization signal (NLS; [51]) found in its N-terminal 48 residues. Despite this, it is evident that a small amount of Tax protein resides in the cytoplasm of mammalian




Fig. 1. Schematic representation of the nuclear and cytoplasmic functions of HTLV-I Tax protein. Tax is shown to activate IKK- and JNK-signaling pathways in the cytoplasm. The IKK-pathway impinges on the nuclear migration of NF- κ B (p50/p65 heterodimer); the JNK-pathway leads to activation of Jun/Fos. In the nucleus, Tax is envisioned to dimerize and to interact with CREB, CBP and P/CAF co-factors as well as with SRF in activating gene transcription. Tax-responsive elements in the HTLV-I LTR reside within the U3.

cells [52]. Thus, in modulating gene expression, Tax is envisioned to have promoter–proximal (i.e. nuclear) as well as promoter–distal (i.e. cytoplasmic) effects (see Fig. 1). Initial studies on Tax demonstrated convincingly that despite its abundant nuclear presence, Tax by itself cannot associate directly with DNA [44]. Contemporaneous findings, however, provided clear evidence that Tax could deregulate the expression of several cellular immediately early (IE) genes. This Tax-mediated dysregulation of gene expression is believed to contribute importantly to its abrogation of normal cellular metabolism [53].

2.1. SRF and AP-1

An early clue as to how HTLV might affect cellular IE genes emerged from the finding that the DNA-binding serum responsive factor (SRF; [54]) could recruit the Tax protein to cellular promoters such as those for *c-fos*, *egr-1* and *egr-2* [47]. Thus, the CC(AT)₆GG motif (CarG box)-binding SRF protein was found to be necessary for Tax's transcriptional activation of a limited subset of promoters [55]; and residues 422–435 of SRF was defined to bind Tax directly [56]. Hence, one mechanistic view is that CarG-box-tethered SRF would bind Tax, thereby bringing the viral oncoprotein to the promoter. The C-terminal activation domain of Tax [57] is then thought to contact directly with TATA-box bound TBP-protein [58] resulting in enhanced transcription.

Many cellular IE genes also contain promoter upstream AP-1 sites. Several studies have shown that HTLV-I transformed T-cells express high levels of AP-1 activity [59,60]. Interestingly, the canonical AP-1 responsive sequence (5'TGACTCA3') bears close resemblance to the three imperfectly conserved core cAMP-responsive palindrome (5' TGACGTCA; [61]) present in the Tax-responsive 21-bp elements found in the HTLV-I LTR. Direct experimental studies have demonstrated that the 21-bp Tax-responsive elements can serve as Jun-responsive sequences [46]. Subsequent work has confirmed that Tax can activate the expression of several cellular promoters through AP-1 sites [62,63]. Despite these reports, currently, it remains unclear whether the major mechanism guiding Tax-activation through AP-1 is via a promoter–proximal enhancement of the DNA-binding activity of AP-1 [64,65], or through a Tax-mediate activation of the JNK-1 kinase [66]. Relevant to the latter possibility, a Tax-binding protein which signals through the TNF- α /AP-1 pathway has recently been described [67].

2.2. CREB, CBP, P/CAF

The HTLV-I LTR (long terminal repeat) contains three imperfectly conserved 21 bp Tax-responsive

(TxRE) sequences, each of which contains a core CREB/ATF binding sites flanked by 5'G- and 3'C-rich residues [29,68]. Highly efficient activation of this viral LTR by Tax is, in part, explained by Tax/CREB/TxRE ternary complex formation at these sites in the LTR [69–73]. However, the ability of Tax to activate transcription via CREB/ATF-sites is context specific, since at other CREB-binding sites (i.e. those found in cellular promoters), Tax–CREB complex formation may occur [70,74,75], but transcriptional activation is not seen.

While the exact details of how Tax activates transcription through CREB/ATF is not completely understood, several findings which contribute important mechanistic insight have been established. It is proposed that the N-terminus of Tax (Fig. 2) directly binds CREB molecules [69,75] docked at CRE-sites in the viral TxRE. Binding of Tax to CREB enhances CREB–CREB homodimerization and heightens resulting association to DNA ([76]; Fig. 3). Because optimal activation through TxREs requires a dimeric form of Tax-protein [77,78], enhanced CREB-dimerization at the HTLV-I LTR could result as a functional consequence of Tax–Tax homodimerization (Fig. 3).

In the Tax/CREB/TxRE ternary complex, Tax makes limited direct contact with DNA [79–81] through its 89–110 amino acid residues ([82]; Fig. 2). Contact with DNA is suggested to result in proper folding of the Tax protein [82] that may lead to a functional presentation of its C-terminal activation domain [57]. Correctly folded Tax protein is known to recruit through its amino acid residues 81–95 ([83]; Fig. 2) the transcriptional co-activators, CBP and p300 ([84–86]; Fig. 3). However, recruitment of CBP/p300 was found to be insufficient for supporting Tax's transcriptional activity [83], suggesting that an additional necessary component for activity remained uncharacterized. This missing factor could be the p300–CBP associated factor, P/CAF, which was recently shown to bind to the C-terminal activation domain of Tax ([87,88]; Figs. 2 and 3). Indeed, Tax mutants that fail to bind to P/CAF correlate with the loss of ability to activate transcription from the HTLV-I LTR [87].

3. Signaling by Tax through NF- κ B

The biology of ATL cells is characterized by increased expression of genes coding for lymphokines [12,19,89] and lymphokine receptors (e.g. IL-2R α ; [19,30–32]). Expression of these genes is in part regulated by the NF- κ B family of transcription factors. The NF- κ B family encompasses several related proteins [90], which can homo- or hetero-dimerize in binding to a GGGRNYYCC DNA-motif. NF- κ B family members function pleiotropically in diverse aspects of immune/inflammatory responses and cellular growth and differ-




Fig. 2. Linear representations of the various functional domains which have been characterized for the 353 amino acid Tax protein. (Top) The N-terminus of Tax contains a nuclear localization signal (NLS) and a zinc-finger (Zn) structure. CREB-binding domain within Tax has also been mapped to the N-terminus. An 'ISGGLC'-sequence has been suggested to contact CREB2 [154]. p300/CBP, P/CAF and DNA-contact domains are indicated. The middle of Tax contains several Leucine (L) residues and is necessary for homodimerization and for NF- κ B activation [155]. At its C-terminus, Tax has an activation domain and a sequence specific to Tax1, but absent from Tax2, which binds to the tumor suppressor hDlg/SAP97 [156]. (Bottom) Schematic representation of single amino acid Tax mutants [94] which were characterized for deficit in transactivation of either the HTLV-I (CREB/ATF pathway) or the HIV-1 (NF- κ B pathway).

entiation [91]. Early HTLV-I research found, interestingly, that Tax activation of genes, such as that for IL-2R α , occurred through NF- κ B binding sites [92,93]. Subsequently, extensive mutagenesis studies [94,95] supported the idea that Tax can generally activate cellular transcription through a NF- κ B-dependent pathway in a manner distinct from its activation of CREB/ATF (Fig. 2). This concept is compatible with the observation that NF- κ B activity is constitutively elevated in primary adult T-cell leukemia cells as well as in Tax-expressing cells [96–100].

3.1. Interaction of Tax and NF- κ B in the nucleus

NF- κ B family members include p105, p100, p65, p52, p50, c-Rel and Rel B [90]. The most frequently observed NF- κ B form is that of a p50–p65 dimer, which is ambiently retained in the cytoplasm by inhibitor ankyrin repeat-containing I κ B molecules (I κ B α , I κ B β , I κ B ϵ , p105, p100 and Bcl-3; reviewed in Ref. [101]). In the commonly accepted paradigm, I κ B α and I κ B β play major roles in sequestering p50–p65 dimer in the cytoplasm. Upon activation of cells by mitogens, cytokines, bacterial lipopolysaccharides, virus infection and/or stress signals, induced-phosphorylation of I κ B α and I κ B β occurs on serines 32 and 36 and serines 19 and 23, respectively (reviewed in Ref. [91]). Phosphorylated I κ B α and I κ B β subunits are then targeted by ubiquitin-ligase component protein, β -TrCP (reviewed in Ref. [102]); this interaction leads to the ubiquitination and proteosomal degradation of the I κ Bs. Upon removal of

I κ Bs, NF- κ B-molecules are freed to migrate from the cytoplasm into the nucleus. Within the nucleus, NF- κ Bs bind promoter upstream DNA-motifs and activate the transcription of a diverse subset of target genes.

Because of Tax's predominant nuclear localization in cells, an initial mechanistic explanation for Tax–NF- κ B interplay invoked events within the nucleus. In this regard, Tax and NF- κ B proteins were observed by immunomicroscopy to colocalize together [103] within nuclear bodies that contain RNA polymerase II and other transcription factors [104]. Tax was also found to capably bind NF- κ B subunits p50 [105] and p52 [106], suggesting a role in direct enhancement of NF- κ B's nuclear transcription function. More recent studies have indicated that the primary action of Tax is to affect NF- κ B signaling in the cytoplasm (see below). However, when cytoplasmic signaling events were inhibited by the use of dominant negative mutants, a small residual activation of NF- κ B by Tax could still be observed [107]. Hence, currently, it cannot be completely excluded that in HTLV-I infected cells nuclear interactive events co-operate with the cytoplasmic effects of Tax to influence overall NF- κ B activation.

3.2. Tax and cytoplasmic signaling of NF- κ B

There are two perspective as to how cytoplasmic Tax modulates NF- κ B activity. Previously, it was known that Tax could directly bind the ankyrin motif contained in molecules such as cyclin-dependent kinase inhibitor p16^{INK4A} [41,108,109]. Such observation sup-




Fig. 3. A model of Tax-mediated transcription through the CREB/ATF pathway. A CREB-dimer is shown to bind to the Tax-responsive elements (TRE) and interact with a Tax homodimer capable of recruiting a molecule(s) of CBP/p300 and P/CAF. This CREB/Tax/CBP/(P/CAF) complex can then influence TATAA-bound TATA-binding protein (TBP) to enhance the initiation of RNA-polymerase II (RNAP II).

ported the reasoning that Tax might physically contact I κ B-molecules through similar ankyrin-motifs, thereby dissociating I κ Bs from cytoplasmically sequestered NF- κ Bs (Fig. 4A). This idea proposes a contact-dependent dissociation mechanism through which Tax would target I κ Bs for proteosomal degradation [110]. Compatible with this view, Tax was found to associate with both I κ B α [111–113] and p105 [114], which can function as a cytoplasmic I κ B-molecule. However, subsequent studies using Tax and I κ B α mutants defined clearly that direct contact between these two molecules insufficiently explains NF- κ B-activation (reviewed in Ref. [45]) suggesting the existence of an alternative activating mechanism(s).

Besides the dissociation model of I κ B from NF- κ B, another general mechanism of NF- κ B activation describes site-specific phosphorylation of I κ B α , followed

by its ubiquitination and degradation [115–117]. A plethora of studies have defined I κ B phosphorylation as a critical regulatory step for NF- κ B activation. Recent insights into how I κ B α phosphorylation occurs inside cells were revealed from extensive characterizations of a 700-kDa complex which contains kinase activity specific for serine 32 and 36 of I κ B α [118,119]. Detailed analyses of this large-kinase-complex resulted in the cloning of two I κ B-kinases — IKK α and IKK β [120–124]. IKK α is an 85-kDa protein, while IKK β is an 87-kDa protein. Both are 52% identical and contain an N-terminal kinase domain, a leucine zipper and a helix-loop-helix motif. Using purified recombinant IKK α and IKK β , it was unambiguously proven that these kinases specifically phosphorylated I κ B α and I κ B β with a slight preference for the former over the latter [125].




Fig. 4. Two models explaining Tax-mediated 'freeing' of NF- κ B from I κ B sequestration. (A) Direct contact of Tax with I κ B leads to dissociation of I κ B from NF- κ B (RelA/p50). (B) Tax is recruited to I κ B/NF- κ B complex through IKK α / β / γ . In this scenario, a Tax-dependent MAP3K activity (not shown) is envisioned to phosphorylate I κ B leading to its ubiquitination and degradation by proteasome.

The large intracellular IKK- α and IKK- β containing-complex, interestingly, was also found to include two members of the MAP kinase kinase kinase (MAP3K) group, MAPK/ERK kinase kinase 1 (MEKK-1) and NF- κ B-inducing kinase (NIK) [126–129]. Very recently, a non-catalytic regulatory subunit for the IKK α and IKK β named IKK γ (or NEMO) was also identified in this complex. cDNAs for human and mouse IKK γ have been cloned [130–132]. Collectively, based on the work of several independent laboratories, it is currently understood that in virtually all settings activation of either IKK α or IKK β requires the IKK γ subunit, since mouse cells deficient in IKK γ fail to phosphorylate and degrade I κ Bs and do not activate NF- κ B in response to diverse stimuli such as TNF α , IL-1, LPS and double-stranded RNA [131]. It is additionally understood that both IKK α and IKK β are activated by phosphorylation on specific serines (S176 and S180 for the former; S177 and S181 for the latter). In this regard, NIK [127,128] and MEKK1 [126,128,129], as well as additional MAP3Ks [133,134] have been invoked as IKK-activating kinases (IKKK). Intriguingly, despite these advances, it remains controversial as to which MAP3K represents the authentic intracellular IKKK needed to activate NF- κ B (reviewed in Refs. [135,136]).

In the context of the IKK-signalosome, how does Tax function mechanistically to activate NF- κ B? Biochemically, over-expressed Tax protein was found by several investigators to be present within the large IKK α -[137–139] and IKK β -[137–140] containing intracellular complex. An initial interpretation was that perhaps Tax contacted either IKK α or IKK β proximally. However, subsequent findings verified that it is IKK γ that binds Tax directly. Thus, IKK γ functionally adapts the Tax-oncoprotein into the large IKK α /IKK β /IKK γ complex [40,141,142].

Simultaneous with the physical characterizations of IKK-Tax association, complementary functional studies have clarified as to how IKK- α and IKK- β activities might be impinged upon by Tax (Fig. 5). First, it was found that dominant negative forms of NIK inhibited Tax-activation of NF- κ B [138,139]. This suggested that Tax acts at a step upstream of NIK-phosphorylation of IKK α / β . Next, work from Gaynor et al. demonstrated a direct association between MEKK1 and Tax, which resulted in a preferential activation of IKK β [140]. A dominant negative form of MEKK1 was shown by these investigators to also inhibit Tax-activation of NF- κ B [140]. Together, these results propose a redundant usage of multiple MAP3Ks by Tax for NF- κ B-activation. Consistent with this redundancy, we have recently observed that over-expression of a dominant negative form of MLK3 (another MAP3K; [134]) further inhibited independently NF- κ B activation by Tax [143]. The molecular interplays between these MAP3Ks




Fig. 5. Summary of sequential events in Tax-activation of nuclear NF- κ B activity. A not yet fully understood interaction occurs between Tax and a MAP3K in the cytoplasm. The identity of the relevant MAP3K remains controversial, although dominant negative forms of NIK, MEKK1 and MLK3 all independently suppress Tax-activation of NF- κ B. Tax is then recruited into the IKK α / β / γ complex. Parenthesis around IKK γ indicates that Tax activation of NF- κ B remains intact in at least one cell line deficient for IKK γ , suggesting that other factors may substitute for IKK γ . The IKK/Tax/MAP3K complex is then postulated to phosphorylate I κ B α . Phosphorylation of I κ B α triggers a ubiquitin-proteosomal degradation pathway which frees NF- κ B (p50/p65) to migrate into the nucleus.

and whether additional MAP3Ks can be utilized by Tax remain to be investigated.

How might one then coalesce extant observations on Tax activation of NF- κ B through the IKK-complex? While several questions remain to be answered, the following statements seemingly describe a plausible functional scenario (Fig. 5). Tax activation of NF- κ B occurs at a point down-stream of the small G proteins and the TNF- α receptor interacting factors [138,140]. At this juncture, Tax bridges the IKK-complex with a MAP3K. Evidence for this bridging mechanism is supported by evidence of direct contact between Tax and both IKK γ [40,141,142] and MEKK1 [140]. MAP3K, as recruited by Tax, is then speculated to phosphorylate IKK α / β leading to a cascade of events (reviewed in Refs. [91,101]) which releases NF- κ B for nuclear migration (Fig. 5). This paradigm is compatible with most of our existing findings on Tax and NF- κ B.

Several important issues regarding NF- κ B activation in HTLV-I-infected cells are, however, not addressed by the above scenario. First, in light of recent findings from knock-out mice that MEKK1 might be dispensable for NF- κ B activation [144], the authentic intracellular MAP3K utilized by Tax requires further definition.

Second, the fact that the gene encoding for IKK γ resides on the X-chromosome [132] suggests IKK- γ functions to be sex-biased [145,146]. Alternatively, there could be additional yet unrecognized adapter proteins that redundantly serve the IKK γ -function. Indeed, in support of this supposition, a B-precursor-cell line, genetically deficient for IKK γ , was previously shown to support fully Tax-activation of NF- κ B [147]. Understanding what are the additional IKK γ -like proteins that could be functionally used by Tax represents a compelling future challenge. Third, additional IKK γ -binding protein [148] which influence NF- κ B-activation has recently been identified. It would be of interest to clarify how such protein(s) might influence the biology of Tax–IKK interaction. Finally, whether Tax alone explains in vivo NF- κ B activation within HTLV-I infected cells has been recently questioned by investigators [100]. The observation that constitutive NF- κ B activation exists in HTLV-I infected cells that do not detectably express Tax suggests that additional virally-encoded (induced) factors should also be carefully investigated [100].

4. Implications for cellular transformation

A critical reason for investigating Tax–NF- κ B interaction is to understand the contribution of this pathway to cellular transformation. Several studies support that Tax activation of NF- κ B contributes to transformation [149–151]. On the other hand, two other studies have suggested that activation through the CREB/ATF pathway by Tax plays a more important role [20,152]. Most recently, experiments using the HTLV-II Tax (Tax2) protein have provided an elegant explanation for these discrepancies. Green et al. demonstrated that NF- κ B-activation by Tax2 provided for initiation of transformation while CREB/ATF-activation served necessarily to maintain the transformed phenotype [153]. An important future goal would be to elucidate how these two pathways cooperate in the important biological process of cellular transformation.

Acknowledgements

I thank Lan Lin for assistance in the preparation of this manuscript and H. Iha, T. Kasai and Y. Iwanaga for critical readings. Due to space limitations, not all relevant papers could be appropriately cited. I apologize to colleagues whose studies may have not been mentioned.

References

- [1] Uchiyama T. Human T cell leukemia virus type I (HTLV-I) and human diseases. *Annu Rev Immunol* 1997;15:15–37.

- [2] Osame M, Usuku K, Izumo S, Ijichi N, Amitani H, Igata A, Matsumoto M, Tara M. HTLV-I associated myelopathy, a new clinical entity. *Lancet* 1986;1(8488):1031–2.
- [3] Poiesz BJ, Ruscetti FW, Gazdar AF, Bunn PA, Minna JD, Gallo RC. Detection and isolation of type C retrovirus particles of fresh and cultured lymphocytes of a patient with cutaneous T-cell lymphoma. *Proc Natl Acad Sci USA* 1980;77:7415–9.
- [4] Kamada N, Sakurai M, Miyamoto K, Sanada I, Sadamori N, Fukuhara S, Abe S, Shiraishi Y, Abe T, Kaneko Y, et al. Chromosome abnormalities in adult T-cell leukemia/lymphoma: a karyotype review committee report. *Cancer Res* 1992;52:1481–93.
- [5] Ueshima Y, Fakuahara S, Hattori T, Hchama T, Takatsuki K, Uchino H. Chromosome studies in adult T-cell leukemia in Japan: significance of trisomy 7. *Blood* 1981;58:420–5.
- [6] Miyamoto K, Sato J, Kitajima KI, Togawa R, Suemaru S, Sanada H, Tanaka T. Adult T-cell leukemia. Chromosome analysis of 15 cases. *Cancer* 1983;52:471–8.
- [7] Rowley JD, Haren JM, Wong-Staal F, Franchini G, Gallo RC, Blattner W. Chromosome pattern in cells from patients positive for human T-cell leukemia/lymphoma virus. In: Gallo RC, Essex ME, Gross L, editors. *Human T-Cell Leukemia/Lymphoma Virus*. New York: Cold Spring Harbor Laboratory, 1984:85–9.
- [8] Whang-Peng J, Bunn PA, Knutsen T, Kao-Shan CS, Broder S, Jaffe ES, Gelmann E, Blattner W, Lofters W, Young RC, Gallo RC. Cytogenetic studies in human T-cell lymphoma virus (HTLV)-positive leukemia-lymphoma in the United States. *J Natl Cancer Inst* 1985;74:357–69.
- [9] Kikuchi M, Mitsui T, Takeshita M, Okamura H, Naitoh H, Eimoto T. Virus associated adult T-cell leukemia (ATL) in Japan: clinical, histological and immunological studies. *Hematol Oncol* 1986;4:67–81.
- [10] Taguchi H, Sonobe H, Yamato K, Takeuchi T, Ookawa K, Kodama H, Ohtsuki Y, Miyoshi I. Central nervous system involvement of adult T-cell leukemia-lymphoma with multinucleated giant cells in the brain, skin, and kidney. *Cancer* 1993;71:133–7.
- [11] Arya SL, Wong-Staal F, Gallo RC. T-cell growth factor gene: Lack of expression in human T-cell leukemia-lymphoma virus-infected cells. *Science* 1984;223:1086–7.
- [12] Arima N, Daitoku Y, Ohgaki S, Fukumori J, Tanaka H, Yamamoto Y, Fujimoto K, Onoue K. Autocrine growth interleukin 2-producing leukemic cells in a patient with adult T cell leukemia. *Blood* 1986;68:779–82.
- [13] Krause BA, Stawhecker JA, Semmes OJ, Jankowski PE, Lweis R, Hinrichs SH. Analysis of potential phosphorylation sites in human T cell leukemia virus type 1 Tax. *J Biomed Sci* 1999;6:206–12.
- [14] Tanaka A, Takahashi C, Yamaoka S, Nosaka T, Maki M, Hatanaka M. Oncogenic transformation by the tax gene of human T-cell leukemia virus type I in vitro. *Proc Natl Acad Sci USA* 1990;87:1071–5.
- [15] Pozzatti R, Vogel J, Jay G. The human T-lymphotropic virus type I tax gene can cooperate with the *ras* oncogene to induce neoplastic transformation of cells. *Mol Cell Biol* 1990;10:413–7.
- [16] Grassmann R, Dengler C, Muller FI, Fleckenstein B, McGuire K, Dokhelar MC, Sodroski JG, Haseltine WA. Transformation to continuous growth of primary human T lymphocytes by human T-cell leukemia virus type I X-region genes transduced by a Herpes virus saimiri vector. *Proc Natl Acad Sci USA* 1989;86:3351–5.
- [17] Grossman WJ, Kimata JT, Wong FH, Zutter M, Ley TJ, Ratner L. Development of leukemia in mice transgenic for the tax gene of human T-cell leukemia virus type I. *Proc Natl Acad Sci USA* 1992;92:1057–61.

- [18] Grassmann R, Berchtold S, Radant I, Alt M, Fleckenstein B, Sodroski JG, Haseltine WA, Ramstedt U. Role of human T-cell leukemia virus type 1 X region proteins in immortalization of primary human lymphocytes in culture. *J Virol* 1992;66:4570–5.
- [19] Inoue J, Seiki M, Taniguchi T, Tsuru S, Yoshida M. Induction of interleukin 2 receptor gene expression by p40x encoded by human T-cell leukemia virus type 1. *EMBO J* 1986;5:2883–8.
- [20] Rosin O, Koch C, Radant I, Schmitt I, Semmes OJ, Jeang KT, Grassman R. A human T-cell leukemia virus Tax variant incapable of activating NF- κ B retains its immortalizing potential for primary T-lymphocytes. *J Biol Chem* 1998;273:6698–703.
- [21] Yamaoka S, Tobe T, Hatanaka M. Tax protein of human T-cell leukemia virus type I is required for maintenance of the transformed phenotype. *Oncogene* 1992;7:433–7.
- [22] Oka T, Sonobe H, Iwata J, Kubonishi I, Satoh H, Takata M, Tanaka Y, Tateno M, Tozawa H, Mori S, Yoshiki T, Ohtsuki Y. Phenotypic progression of a rat lymphoid cell line immortalized by human T-lymphotropic virus type 1 to induce lymphoma/leukemia-like disease in rats. *J Virol* 1992;66:6686–94.
- [23] Nerenberg M, Hinrichs SH, Reynolds RK, Khoury G, Jay G. The tat gene of human T-lymphotropic virus type 1 induces mesenchymal tumors in transgenic mice. *Science* 1987;237:1324–9.
- [24] Green JE, Hinrichs SH, Vogel J, Jay G. Exocrinopathy resembling Sjogren's syndrome in HTLV-I tax transgenic mice. *Nature* 1989;341:72–4.
- [25] Peebles RS, Maliszewski CR, Sato TA, Hanley HJ, Maroulakou IG, Hunziker R, Schneck JP, Green JE. Abnormal B-cell function in HTLV-I-tax transgenic mice. *Oncogene* 1995;10:1045–51.
- [26] Grossman WJ, Kimata JT, Wong FH, Zutter M, Ley TJ, Ratner L. Development of leukemia in mice transgenic for the tax gene of human T-cell leukemia virus type I. *Proc Natl Acad Sci USA* 1995;92:1057–61.
- [27] Ross TM, Pettiford SM, Green PL. The tax gene of human T-cell leukemia virus type 2 is essential for transformation of human T lymphocytes. *J Virol* 1996;70:5194–202.
- [28] Seiki M, Inoue J, Takeda T, Yoshida M. Direct evidence that p40x of human T-cell leukemia virus type I trans-acting transcriptional activator. *EMBO J* 1986;5:561–5.
- [29] Brady J, Jeang KT, Duvall J, Khoury G. Identification of p40x-responsive regulatory sequences within the human T-cell leukemia virus type I long terminal repeat. *J Virol* 1987;61:2175–81.
- [30] Maruyama M, Shibuya H, Harada H, Hatakeyama M, Seiki M, Fujita T, Inoue J, Yoshida M, Taniguchi T. Evidence for aberrant activation of the interleukin-2 autocrine loop by HTLV-I-encoded p40x and T3/Ti complex triggering. *Cell* 1987;48:343–50.
- [31] Cross SL, Feinberg MB, Wolf JB, Holbrook NJ, Wong-Staal F, Leonard WJ. Regulation of the human interleukin-2 receptor alpha chain promoter: activation of a nonfunctional promoter by the transactivator gene of HTLV-I. *Cell* 1987;49:47–56.
- [32] Siekevitz M, Feinberg MB, Holbrook N, Wong-Staal F, Greene WC. Activation of interleukin 2 and interleukin 2 receptor (Tac) promoter expression by the *trans*-activator (tat) gene product of human T-cell leukemia virus, type I. *Proc Natl Acad Sci USA* 1987;85:5389–93.
- [33] Fujii M, Sassone-Corsi P, Verma IM. c-fos promoter *trans*-activation by the tax1 protein of human T-cell leukemia virus type I. *Proc Natl Acad Sci USA* 1988;85:8526–30.
- [34] Wano Y, Feinberg M, Hosking JB, Bogerg H, Greene WC. Stable expression of the tax gene of type I human T-cell leukemia virus in human T cells activates specific cellular genes involved in growth. *Proc Natl Acad Sci USA* 1988;85:9733–7.
- [35] Nagata K, Ohtani K, Nakamura M, Sagamura K. Activation of endogenous c-fos proto-oncogene expression by human T-cell leukemia virus type I-encoded p40^{tax} protein in the human T-cell line. *J Virol* 1989;63:3220–6.
- [36] Hollsberg P. Mechanisms of T-cell activation by human T-cell lymphotropic virus type I. *Micro Mol Biol Rev* 1999;63:308–33.
- [37] Miyatake S, Seiki M, Yoshida M, Arai K. T-cell activation signals and human T-cell leukemia virus type-I-encoded p40x protein activate the mouse granulocyte-macrophage colony-stimulating factor gene through a common DNA element. *Mol Cell Biol* 1988;8:5581–7.
- [38] Miyoshi I, Kubonishi I, Sumida M, Hiraki S, Tsubota T, Kimura I, Miyamoto K, Sato J. A novel T-cell line derived from adult T-cell leukemia. *Gann* 1980;71:155–6.
- [39] Colgin MA, Nyborg J. The human T-cell leukemia virus type 1 oncoprotein tax inhibits the transcriptional activity of c-myc through competition for the CREB binding protein. *J Virol* 1998;72:9396–9.
- [40] Jin DY, Giordano V, Kibler KV, Nakano H, Jeang KT. Human T-cell leukemia virus type I Tax interacts directly with I κ B kinase γ . *J Biol Chem* 1999;274:17402–5.
- [41] Low KG, Dorner LF, Fernando DB, Grossman J, Jeang KT, Comb MJ. Human T-cell leukemia virus type 1 Tax releases cell cycle arrest induced by p16INK4a. *J Virol* 1997;71:1956–62.
- [42] Ohtani K, Nakamura M, Saito S, Nagata K, Sugamura K, Hinuma Y. Electroporation: application to human lymphoid cell lines for stable introduction of a transactivator gene of human T-cell leukemia virus type I. *Nucleic Acids Res* 1989;17:1589–604.
- [43] Van Orden K, Giebler HA, Lemasson L, Gonzales M, Nyborg JK. Binding of p53 to the KIX domain of CREB binding protein. *J Biol Chem* 1999;274:26321–8.
- [44] Franklin AA, Nyborg JK. Mechanisms of Tax regulation of human T cell leukemia virus type I gene expression. *J Biomed Sci* 1995;2:17–29.
- [45] Sun SC, Ballard DW. Persistent activation of NF- κ B by the Tax transforming protein of HTLV-I: hijacking cellular I κ B kinases. *Oncogene* 1999;18:6948–58.
- [46] Jeang KT, Chiu R, Santos E, Kim SJ. Induction of the HTLV-I LTR by jun occurs through the Tax-responsive 21-bp elements. *Virology* 1991;181:218–27.
- [47] Fujii M, Tsuchiya J, Chuhjo T, Akizawa T, Seiki M. Interaction of HTLV-I Tax1 with p67SRF causes the aberrant induction of cellular immediate early genes through CAR γ boxes. *Genes Dev* 1992;6:2066–76.
- [48] Hiscott J, Petropoulos L, Lacoste J. Molecular interactions between HTLV-I protein and the NF- κ B/I κ B transcription complex. *Virology* 1995;214:3–11.
- [49] Li XH, Gaynor RB. Regulation of NF- κ B by the HTLV-I tax protein. *Gene Exp* 1999;7:233–45.
- [50] Neuveut C, Jin DY, Semmes OJ, Diella F, Callahan R, Jeang KT. Divergent subcellular locations of HTLV-I Tax and Int-6: A contrast between in vitro protein–protein binding and intracellular protein co-localization. *J Biomed Sci* 1997;4:229–34.
- [51] Smith MR, Greene WC. Characterization of a novel nuclear localization signal in the HTLV-I tax transactivator protein. *Virology* 1992;187:316–20.
- [52] Nicot C, Tie F, Giam CZ. Cytoplasmic forms of human T-cell leukemia virus type I Tax induce NF- κ B activation. *J Virol* 1998;72:6777–84.
- [53] Yoshida M. HTLV-I Tax: regulation of gene expression and disease. *Trends Microbiol* 1993;1:131–5.
- [54] Herdegen T, Leah JD. Inducible and constitutive transcription factors in the mammalian nervous system: control of gene expression by Jun, Fos and Krox and CREB/ATF proteins. *Brain Res Brain Res Rev* 1998;28:370–490.

- [55] Fujii M, Tsuchiya J, Chuhjo T, Minamino T, Miyamoto K, Seiki M. Serum response factor has functional roles both in indirect bind to the CARG box and in the transcriptional activation function human T-cell leukemia virus type I Tax. *J Virol* 1994;68:7275–83.
- [56] Fujii M, Chuhjo T, Minamino T, Masaaki N, Miyamoto K, Seiki M. Identification of the Tax interaction region of serum response factor that mediates the aberrant induction of immediate early genes through CARG boxes by HTLV-I Tax. *Oncogene* 1995;11:7–14.
- [57] Semmes OJ, Jeang KT. Definition of a minimal activation domain in HTLV-I Tax. *J Virol* 1995;69:1827–33.
- [58] Caron C, Rousset R, Beraud C, Moncollin V, Egly JM, Jalinot P. Functional and biochemical interaction of the HTLV-I Tax1 transactivator with TBP. *EMBO J* 1993;12:4269–78.
- [59] Mulloy JC, Kislyakova T, Cereseto A, Casareto L, LoMonico A, Fullen J, Lorenzi MV, Cara A, Nicot C, Giam C, Franchini G. Human T-cell lymphotropic/leukemia virus type I Tax abrogates p53-induced cell cycle arrest and apoptosis through its CREB/ATF functional domain. *J Virol* 1998;72:8852–60.
- [60] Fujii M, Niki T, Mori T, Matsuda T, Matsui M, Nomura N, Seiki M. HTLV-I Tax induces expression of various immediate early serum responsive genes. *Oncogene* 1991;6:1023–9.
- [61] Jeang KT, Boros I, Brady J, Radonovich M, Khoury G. Characterization of cellular factors that interact with the human T-cell leukemia virus type I p40x-responsive 21-base-pair sequence. *J Virol* 1988;62:499–509.
- [62] Ejima E, Rosenblatt JD, Massari M, Quan E, Stephens D, Rosen CA. Cell-type-specific transactivation of the parathyroid hormone-related protein gene promoter by the human T-cell leukemia vi type I (HTLV-I) tax and HTLV-II tax proteins. *Blood* 1993;81:1017–24.
- [63] Liu XD, Chen XL, Zachar V, Chang CS, Ebbesen P. Transcriptional activation of human TR3/Nur77 gene expression by human T-lymphotropic virus type I Tax protein through two AP-1-like elements. *J Gen Virol* 1999;80:3073–81.
- [64] Armstrong AP, Franklin AA, Uittenbogaard MN, Giebler HA, Nyborg A. Pleiotropic effect of the human T-cell leukemia virus Tax protein on the DNA binding activity of eukaryotic transcription factors. *Proc Natl Acad Sci USA* 1993;90:7303–7.
- [65] Mori N, Fujii M, Iwai K, Ikeda S, Yamasaki Y, Jata T, Yamada Y, Tan Y, Tomonaga M, Yamamoto N. Constitutive activation of transcription factor AP-1 in primary adult T-cell leukemia cells. *Blood* 2000;95:3915–21.
- [66] Jin DY, Teramoto H, Giam CZ, Chun RF, Gutkind S, Jeang KT. A human suppressor of c-Jun N-terminal kinase 1 activation by tumor necrosis factor α . *J Biol Chem* 1997;272:25816–23.
- [67] Iha H, Kasai T, Kibler KV, Iwanaga Y, Tsurugi K, Jeang KT. Pleiotropic effects of HTLV Type I Tax protein on cellular metabolism: mitotic checkpoint abrogation and NF- κ B activation. *AIDS Res Hum Retroviruses* 2000;16(16):1633–8.
- [68] Jeang KT, Boros I, Brady J, Radonovich M, Khoury G. Identification of cellular factors that bind to the 21bp p40x responsive region in the HTLV-I LTR. *J Virol* 1988;62:4499–509.
- [69] Goren I, Semmes OJ, Jeang KT, Moelling K. The amino terminus of tax is required for interaction with the cyclic AMP response element binding protein. *J Virol* 1995;69:5806–11.
- [70] Laurance ME, Kwok RPS, Huang MS, Richards JP, Lundblad JR, Goodman RH. Differential activation of viral and cellular promoters by human T-cell lymphotropic virus-I Tax and cAMP-responsive element modulator isoforms. *J Biol Chem* 1997;272:2646–51.
- [71] Paca-Uccaralertkun S, Zhao LJ, Adya N, Cross JV, Cullen BR, Boros IM, Giam CZ. In vitro selection of DNA elements highly responsive to the human T-cell lymphotropic virus type I transcriptional activator, Tax. *Mol Cell Biol* 1994;14:456–62.
- [72] Brauweiller A, Garl P, Franklin AA, Giebler HA, Nyborg JK. A molecular mechanism for human T-cell leukemia virus latency and Tax transactivation. *J Biol Chem* 1996;270:12814–22.
- [73] Yin MJ, Gaynor RB. Complex formation between CREB and Tax enhances the bind affinity of CREB for the human T-cell leukemia virus types 1 21 base-pair repeats. *Mol Cell Biol* 1996;16:3156–68.
- [74] Suzuki T, Fujisawa JI, Toita M, Yoshida M. The trans-activator Tax of human T-cell leukemia virus type I (HTLV-I) interacts with cAMP-responsive element (CRE) binding and CRE modulator proteins that bind to the 21-base-pair enhancer of HTLV-I. *Proc Natl Acad Sci USA* 1993;90:610–4.
- [75] Adya N, Giam CZ. Distinct regions in human T-cell lymphotropic virus type I Tax mediate interactions with activator protein CREB and basal transcription factors. *J Virol* 1995;69:1834–41.
- [76] Baranger AM, Palmer CR, Hamm MK, Giebler HA, Brauweiler A, Nyborg JK, Schepartz A. Mechanism of DNA-binding enhancement by the human T-cell leukaemia virus transactivator, tax. *Nature* 1995;376:606–8.
- [77] Jin DY, Jeang KT. HTLV-I tax self-association in optimal trans-activation function. *Nucleic Acids Res* 1997;25:379–87.
- [78] Tie F, Adya N, Greene WC, Giam CZ. Interaction of the human T-lymphotropic virus type I Tax dimer with CREB and the viral 21-base-pair repeat. *J Virol* 1996;70:8368–74.
- [79] Lenzmeier BA, Giebler HA, Nyborg JK. Human T-cell leukemia virus type I Tax requires direct access to DNA for recruitment of CREB binding protein to the viral promoter. *Mol Cell Biol* 1998;18:721–31.
- [80] Lundblad JR, Kwok RP, Laurance ME, Huang MS, Richard JP, Brennan RG, Goodman RH. The human T-cell leukemia virus-1 transcriptional activator Tax enhances cAMP-responsive element-binding protein (CREB) binding activity through interactions with the DNA minor groove. *J Biol Chem* 1998;273:19251–9.
- [81] Kimzey AL, Dynan WS. Specific regions of contact between human T-cell leukemia virus type I Tax protein and DNA identified by photocross-linking. *J Biol Chem* 1998;273:13768–75.
- [82] Kimzey AL, Dynan WS. Identification of a human T-cell leukemia virus type I Tax peptide contact with DNA. *J Biol Chem* 1999;274:34226–32.
- [83] Harrod R, Tang Y, Nicot C, Lu HS, Vassilev A, Nakatani Y, Giam CZ. An exposed KID-like domain in human T-cell lymphotropic virus type I Tax is responsible for the recruitment of coactivators CBP/p300. *Mol Cell Biol* 1998;18:5052–61.
- [84] Kwok RP, Laurance ME, Lundblad JR, Goldman PS, Shih H, Connor LM, Marriott SJ, Goodman RH. Control of cAMP-regulated enhancers by the viral transactivator Tax through CREB and the co-activator CBP. *Nature* 1996;380:642–6.
- [85] Giebler HA, Loring JE, Van Orden K, Colgin MA, Garrus JE, Escudero KW, Brauweiler A, Nyborg JK. Anchoring of CREB binding protein to the human T-cell leukemia virus type I promoter: a molecular mechanism of Tax transactivation. *Mol Cell Biol* 1997;17:5156–64.
- [86] Bex F, Yin MJ, Burny A, Gaynor RB. Differential transcriptional activation by human T-cell leukemia virus type I Tax mutants is mediated by distinct interactions with CREB binding protein and p300. *Mol Cell Biol* 1998;18:2392–405.
- [87] Harrod R, Kuo YL, Tang Y, Yao Y, Vassiev A, Nakatani Y, Giam CZ. p300 and p300/cAMP-responsive element-binding protein associated factor interact with human T-cell lymphotropic virus types-1 Tax multi-histone acetyltransferase/activator-enhancer complex. *J Biol Chem* 2000;275:11852–7.
- [88] Jiang H, Lu H, Schilz RL, Pise-Masison CA, Ogryzko VV, Nakatani Y, Brady JN. P/CAF interacts with tax and stimulates tax transactivation in a histone acetyltransferase-independent manner. *Mol Cell Biol* 1999;19:8136–45.

- [89] Miyatake S, Seiki M, Malefijt RD, Heike T, Fujisawa J, Takebe Y, Nishida J, Shilomai J, Yokota T, Yoshida M, Arai K, Arai N. Activation of T-cell derived lymphokine genes in T-cells and fibroblasts: Effects of human T-cell leukemia virus type I, p40 protein and bovine papilloma virus encoded E2 protein. *Nucleic Acids Res* 1988;16:6547–66.
- [90] DeLuca C, Kwon H, Lin R, Wainberg M, Hiscott J. NF- κ B activation and HIV-1 induced apoptosis. *Cyto Growth Factor Rev* 1999;10:235–53.
- [91] Baldwin AS Jr. The NF-kappa B and I kappa B proteins: new discoveries and insights. *Annu Rev Immunol* 1996;14:649–83.
- [92] Leung K, Nabel GJ. HTLV-I transactivator induces interleukin-2 receptor expression through an NF- κ B-like factor. *Nature* 1988;333:776–8.
- [93] Lowenthal JW, Bohnlein E, Ballard DW, Greene WC. Regulation of interleukin 2 receptor α subunit (Tac or CD25 antigen) gene expression: Binding of inducible nuclear proteins to discrete promoter sequences correlates with transcriptional activation. *Proc Natl Acad Sci USA* 1988;85:4468–73.
- [94] Semmes OJ, Jeang KT. Mutational analysis of human T-cell leukemia virus type I Tax: regions necessary for function determined with 47 mutant proteins. *J Virol* 1992;66:7183–92.
- [95] Smith MR, Greene WC. Identification of HTLV-I Tax transactivator mutants exhibiting novel transcriptional phenotypes. *Genes Dev* 1990;4:1875–85.
- [96] Kanno T, Broen K, Serbenlist U. Evidence in support of a role for human T-cell leukemia virus type I Tax in activating NF- κ B via stimulation of signaling pathways. *J Biol Chem* 1995;270:11745–8.
- [97] Lacoste J, Petropoulos L, Pepin N, Hiscott J. Constitutive phosphorylation and turnover of I κ B α in human T-cell leukemia virus type I-infected and Tax-expressing T cells. *J Virol* 1995;69:864–9.
- [98] Maggirwar SB, Harhaj E, Sun SC. Activation of NF- κ B/Rel by Tax involves degradation I κ B α and blocked by a proteasome inhibitor. *Oncogene* 1995;11:993–8.
- [99] McKinsey TA, Brockman JA, Scherer DC, Al-Murrani SW, Green PL, Ballard DW. Inactivation of I κ B α by the Tax protein of human T-cell leukemia virus type 1: A potential mechanism for constitutive induction of NF- κ B. *Mol Cell Biol* 1996;16:2083–90.
- [100] Mori N, Fujii M, Ikeda S, Yamada Y, Tomonaga M, Ballard DW, Yamamoto N. Constitutive activation of NF-kappaB in primary adult T-cell leukemia cells. *Blood* 1999;93:2360–8.
- [101] Verma IM, Stevenson JK, Schwarz EM, Van Antwerp D, Miyamoto S. Rel/NF-kappa B/I kappa B family: intimate tales of association and dissociation. *Genes Dev* 1995;9:2723–35.
- [102] Maniatis T. A ubiquitin ligase complex essential for the NF-kappaB, Wnt/Wingless, and Hedgehog signaling pathways. *Gene Dev* 1999;13:505–10.
- [103] Bex F, McDowall A, Burny A, Gaynor R. The human T-cell leukemia virus type I transactivator protein Tax colocalizes in unique nuclear structures with NF- κ B proteins. *J Virol* 1997;71:3484–97.
- [104] Semmes OJ, Jeang KT. Localization of human T-cell leukemia virus type I Tax to subnuclear compartments that overlap with interchromatin speckles. *J Virol* 1996;70:6347–57.
- [105] Suzuki T, Hirai H, Fujisawa J, Fujita T, Yoshida M. A transactivator Tax of human T-cell leukemia virus type 1 binds to NF- κ B p50 and serum response factor (SRF) and associates with enhancer DNAs of the NF- κ B site and CArG box. *Oncogene* 1993;8:2391–7.
- [106] Murakami T, Hirai H, Suzuki T, Fujisawa JI, Yoshida M. HTLV-I Tax enhances NF- κ B2 expression and binds to the products p52 and p100, but does not suppress the inhibitory function of p100. *Virology* 1995;206:1066–74.
- [107] Uhlik M, Good LF, Xiao G, Harhaj EW, Zandi E, Karin M, Sun SC. NF- κ B-inducing kinase and I κ B kinase participate in human T-cell leukemia virus I Tax-mediated NF- κ B activation. *J Biol Chem* 1998;273:21132–6.
- [108] Suzuki T, Kitao S, Matsushime H, Yoshida M. HTLV-I Tax protein interacts with cyclin-dependent kinase inhibitor p16INK4A and counteracts its inhibitory activity towards CDK4. *EMBO J* 1996;15:1607–14.
- [109] Neuveut C, Low KG, Maldarelli F, Schmitt I, Majone F, Grassmann R, Jeang KT. Human T-cell leukemia virus type I Tax and cell cycle progression: role of Cyclin D-cdk and p110Rb. *Mol Cell Biol* 1998;18:3620–32.
- [110] Beraud C, Greene WC. Interaction of HTLV-I tax with the human proteasome: implications for NF-kappa B induction. *J Acq Immune Defic Syndr Hum Retrovirol* 1996;1(13):S76–84.
- [111] Hirai H, Suzuki T, Fujisawa J, Inoue J, Yoshida M. Tax protein of human T-cell leukemia virus type I binds to the ankyrin motifs of inhibitory factor kappa B and induces nuclear translocation of transcription factor NF-kappa B proteins for transcriptional activation. *Proc Natl Acad Sci USA* 1994;91:3584–8.
- [112] Suzuki T, Hirai H, Murakami T, Yoshida M. Tax protein of HTLV-I destabilizes the complexes of NF-kappa-B and I kappa B-alpha and induces nuclear translocation of NF- κ B for transcriptional activation. *Oncogene* 1995;10:1199–207.
- [113] Petropoulos L, Hiscott J. Association between HTLV-I Tax and I κ B α is dependent on the I κ B α phosphorylation state. *Virology* 1998;252:189–99.
- [114] Hirai H, Fujisawa J, Suzuki T, Ueda K, Muramatsu M, Tsuboi A, Arai, Yoshida M. Transcriptional activator Tax of HTLV-I binds to the NF-kappa B precursor p105. *Oncogene* 1992;7:1737–42.
- [115] Palombella VJ, Rando OJ, Goldberg AL, Maniatis T. The ubiquitin-proteasome pathway is required for processing the NF-kappa B1 precursor protein and the activation of NF-kappa-B. *Cell* 1994;78:773–85.
- [116] Brown K, Gerstberger S, Carlson L, Franzoso G, Siebenlist U. Control of I kappa B-alpha proteolysis by site-specific, signal-induced phosphorylation. *Science* 1995;267:1485–8.
- [117] Traenckner EB, Pahl HL, Henkel T, Schmidt KN, Wilk S, Baeuerle PA. Phosphorylation of human I kappa B-alpha on serines 32 and 36 control I kappa B-alpha proteolysis and NF-kappa B activation in response to diverse stimuli. *EMBO J* 1995;14:2876–83.
- [118] Chen ZJ, Parent L, Maniatis T. Site-specific phosphorylation of I κ B α by a novel ubiquitination-dependent protein kinase activity. *Cell* 1996;84:853–62.
- [119] Lee FS, Hagler J, Chen ZJ, Maniatis T. Activation of the I κ B α kinase complex by MEKK1, a kinase of the JNK pathway. *Cell* 1997;88:213–22.
- [120] DiDonato JA, Hayakawa M, Rothwarf DM, Zandi E, Karin M. A cytokine-responsive I κ B kinase that activates the transcription factor NF- κ B. *Nature* 1997;388:548–54.
- [121] Mercurio F, Zhu H, Murray BW, Shevchenko A, Bennett BL, Li J, Young DB, Barbosa M, Mann M. IKK-1 and IKK-2: cytokine-activated I κ B kinases essential for NF- κ B activation. *Science* 1997;278:860–6.
- [122] Zandi E, Rothwarf DM, Delhase M, Hayakawa M, Karin M. The I κ B kinase complex (IKK) contains two kinase sub-units, IKK α and IKK β , necessary for I κ B phosphorylation and NF- κ B activation. *Cell* 1997;91:243–52.
- [123] Regnier CH, Song HY, Gao X, Goeddel DV, Cao Z, Rothe M. Identification and characterization of an I κ B kinase. *Cell* 1997;90:373–83.
- [124] Woronicz JD, Gao X, Cao Z, Rothe M, Goeddel DV. I κ B kinase- β : NF- κ B activation and complex formation with I κ B kinase- α and NIK. *Science* 1997;278:866–9.

- [125] Zandi E, Chen Y, Karin M. Direct phosphorylation of I κ B α by IKK α and IKK β : discrimination between free and NF- κ B-bound substrate. *Science* 1998;281:1360–3.
- [126] Lee FS, Peters R, Dang LC, Maniatis T. MEKK1 activates both I κ B kinase α and I κ B kinase β . *Proc Natl Acad Sci USA* 1998;95:9319–24.
- [127] Ling L, Cao Z, Goeddel DV. NF- κ B-inducing kinase activates IKK- α by phosphorylation of Ser-176. *Proc Natl Acad Sci USA* 1998;95:3792–97.
- [128] Nakano H, Shindo M, Sakon S, Nishinaka S, Mihara M, Yagita H, Okumura K. Differential regulation of I κ B kinase α and β by two upstream kinase, NF- κ B-inducing kinase and mitogen-activated protein kinase/ERK kinase kinase-1. *Proc Natl Acad Sci USA* 1998;95:3537–42.
- [129] Nemoto S, DiDinatto JA, Lin A. Coordinate regulation of I κ B kinases by mitogen-activated protein kinase kinase 1 and NF- κ B-inducing kinase. *Mol Cell Biol* 1998;18:7336–43.
- [130] Rothwarf DM, Zandi E, Natoli G, Karin M. IKK- γ is an essential regulatory subunit of the I κ B kinase complex. *Nature* 1998;395:297–300.
- [131] Yamaoka S, Courtois G, Bessia C, Whiteside ST, Weil R, Agou F, Kirk HE, Kay RJ, Israel A. Complementation cloning of NEMO, a component of the I κ B kinase complex essential for NF- κ B activation. *Cell* 1998;93:1231–40.
- [132] Jin DY, Jeang KT. Isolation of full-length cDNA and chromosomal localization of human NF- κ B modulator NEMO to Xq28. *J Biomed Sci* 1999;6:115–20.
- [133] Zhao Q, Lee FS. Mitogen-activated protein kinase/ERK kinase kinases 2 and 3 activate nuclear factor- κ B through I κ B kinase- α and I κ B kinase- β . *J Biol Chem* 1999;274:8355–8.
- [134] Hehner SP, Hofmann TG, Ushmorov A, Dienz O, Leung IW, Lassam N, Scheidereit C, Droge W, Schmitz ML. Mixed-lineage kinase 3 delivers CD3/CD28-derived signals into the I κ B kinase complex. *Mol Cell Biol* 2000;20:2556–68.
- [135] Karin M, Delhase M. JNK or IKK, AP-1 or NF- κ B, which are the targets for MEK kinase 1 action? *Proc Natl Acad Sci USA* 1998;95:9067–9.
- [136] Karin M, Delhase M. The I κ B kinase (IKK) and NF- κ B: key elements of proinflammatory signalling. *Immunology* 2000;12:85–98.
- [137] Chu ZL, DiDonato JA, Hawiger J, Ballard DW. The Tax oncoprotein of human T-cell leukemia virus type 1 associates with and persistently activates I κ B kinases containing IKK α and IKK β . *J Biol Chem* 1998;273:15891–4.
- [138] Geleziunas R, Ferrell S, Lin X, Mu Y, Cuninghame ET Jr, Grant M, Connelly MA, Hambor JE, Marcu KB, Greene WC. Human T-cell leukemia virus type 1 tax induction of NF- κ B involves activation of the I κ B kinase α (IKK α) and IKK β cellular kinases. *Mol Cell Biol* 1998;18:5157–65.
- [139] Uhlik M, Good KF, Xiao G, Harhaj EW, Zandi E, Karin M, Sun SC. NF- κ B-inducing kinase and I κ B kinase participate in human T-cell leukemia virus I Tax-mediated NF- κ B activation. *J Biol Chem* 1998;273:21132–6.
- [140] Jin MJ, Christerson LB, Yamamoto Y, Kwak YT, Xu S, Mercurio F, Barbosa M, Cobb MH, Gaynor RB. HTLV-I Tax protein binds to MEKK1 to stimulate I κ B kinase activity and NF- κ B activation. *Cell* 1998;93:875–84.
- [141] Chu ZL, Shin YA, Yang JM, DiDonato JA, Ballard DW. IKK γ mediates the interaction of cellular I κ B kinase with the tax transforming protein of human T cell leukemia virus type 1. *J Biol Chem* 1999;274:15297–300.
- [142] Harhaj EW, Sun SC. IKK γ serves as a docking subunit of the I κ B kinase (IKK) and mediates interaction of IKK with the human T-cell leukemia virus Tax protein. *J Biol Chem* 1999;274:22911–4.
- [143] Iha H, Jeang KT. Unpublished observation.
- [144] Yuiri T, Ware M, Widmann C, Oyer R, Russell D, Chan E, Zaitzu Y, Clarke P, Tyler K, Oka Y, Fanger GR, Henson P, Johnson GL. MEK kinase 1 gene disruption alters cell migration and c-Jun NH $_2$ -terminal kinase regulation but does not cause a measurable defect in NF- κ B activation. *PNAS* 2000;13:7272–7.
- [145] Makris C, Godfrey VL, Krahn-Senfleiben G, Takahashi T, Roberts JL, Schwarz T, Feng L, Johnson RS, Karin M. Female mice heterozygous for IKK γ /NEMO deficiencies develop a dermatopathy similar to the human X-linked disorder incontinentia pigmenti. *Mol Cell* 2000;5:969–79.
- [146] Smahi A, Courtois G, Vabres P, Yamaoka S, Heuertz S, et al. Genomic rearrangement in NEMO impairs NF- κ B activation and is a cause of incontinentia pigmenti. *Nature* 2000;405:466–72.
- [147] Courtois G, Whiteside ST, Sibley CH, Israel A. Characterization of a mutant cell line that does not activate NF- κ B in response to multiple stimuli. *Mol Cell Biol* 1997;17:1441–9.
- [148] Li S, Commane M, Nie H, Hua X, Chatterjee-Kishore M, Wald D, Haag M, Stark GR. Act1, an NF- κ B-activating protein. *PNAS* 2000;97:10489–93.
- [149] Roberk MD, Ratner L. immortalization of CD4 $^{+}$ and CD8 $^{+}$ T lymphocytes by human T-cell leukemia virus type 1 Tax mutants expressed in a functional molecular clone. *J Virol* 1999;73:4856–65.
- [150] Matsumoto K, Shibata H, Fujisawa J, Inoue H, Hakura A, Tsukahara T, Fujii M. Human T-cell leukemia virus type 1 Tax protein transforms rat fibroblasts via two distinct pathways. *J Virol* 1997;71:4445–51.
- [151] Yamaoka S, Inoue H, Sakurai M, Sugiyama T, Hazama M, Yamada T, Hatanaka M. Constitutive activation of NF- κ B is essential for transformation of rat fibroblasts by the human T-cell leukemia virus type I Tax protein. *EMBO J* 1996;15:873–87.
- [152] Smith MR, Greene WC. Type I human T cell leukemia virus tax protein transforms rat fibroblasts through the cyclic adenosine monophosphate responsive element binding protein/activating transcription factor pathway. *J Clin Invest* 1991;88:1038–42.
- [153] Ross TM, Narayan M, Fang ZY, Minella AC, Green P. Human T-cell leukemia virus type 2 Tax mutants that selectively abrogate NF κ B or CREB/ATF activation fail to transform primary human T cells. *J Virol* 2000;74:2655–62.
- [154] Gachon F, Peleraux A, Thebault S, Dick J, Lemasson I, Devaux C, Mesnard JM. CREB-2, a cellular CRE-dependent transcription repressor functions in association with Tax as an activator of the human T-cell leukemia virus type I promoter. *J Virol* 1998;72:8332–7.
- [155] Kuo YL, Tang Y, Harrod R, Cai P, Giam CZ. The KID-like region of the HTLV-1 Tax is important for NF- κ B activation. *AIDS Res Hum Retroviruses* 2000;16(16):1607–12.
- [156] Suzuki T, Ohsugi Y, Uchida-Toita M, Akiyama T, Yoshida M. Tax oncoprotein of HTLV-1 binds to the human homologue of Drosophila discs large tumor suppressor protein, hDLG, and perturbs its function in cell growth control. *Oncogene* 1999;18:5967–72.